

WASL rabbit pAb

Cat No.: ES11262

For research use only

Overview

Product Name WASL rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human;Rat;Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein . at

AA range: 190-270

Specificity WASL Polyclonal Antibody detects endogenous

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

StorageStore at -20°C. Avoid repeated freeze-thaw cycles.Protein NameNeural Wiskott-Aldrich syndrome protein (N-WASP)

Gene Name WASL

Cellular localization Cytoplasm, cytoskeleton . Nucleus . Cytoplasm .

Preferentially localized in the cytoplasm when phosphorylated and in the nucleus when

unphosphorylated (By similarity). Exported from the nucleus by an nuclear export signal (NES)-dependent

mechanism to the cytoplasm (By similarity).

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 55kD
Human Gene ID 8976
Human Swiss-Prot Number 000401

Alternative Names

Background This gene encodes a member of the Wiskott-Aldrich

syndrome (WAS) protein family. Wiskott-Aldrich syndrome proteins share similar domain structure, and associate with a variety of signaling molecules



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



to alter the actin cytoskeleton. The encoded protein is highly expressed in neural tissues, and interacts with several proteins involved in cytoskeletal organization, including cell division control protein 42 (CDC42) and the actin-related protein-2/3 (ARP2/3) complex. The encoded protein may be involved in the formation of long actin microspikes, and in neurite extension. [provided by RefSeq, Jul 2013],

