

TKNK rabbit pAb

Cat No.: ES12686

For research use only

Overview

Product Name TKNK rabbit pAb

Host species Rabbit

Applications WB;ELISA;IHC Species Cross-Reactivity Human; Mouse;Rat

Recommended dilutions WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000

Immunogen Synthesized peptide derived from human TKNK AA

Specificity This antibody detects endogenous levels of TKNK at

range: 6-56

Human/Mouse/Rat

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name TKNK

Gene Name TAC3 NKNB UNQ585/PRO1155

Cellular localization Secreted.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

Observed band

Human Gene ID 6866 Human Swiss-Prot Number Q9UHF0

Alternative Names

Background This gene encodes a member of the tachykinin

family of secreted neuropeptides. The encoded preproprotein is proteolytically processed to generate the mature peptide, which is primarily expressed in the central and peripheral nervous systems and functions as a neurotransmitter. This peptide is the ligand for the neurokinin-3 receptor.

This protein is also expressed in the outer

syncytiotrophoblast of the placenta and may be



+86-27-59760950


ELKbio@ELKbiotech.com

www.elkbiotech.com



associated with pregnancy-induced hypertension and pre-eclampsia. Mutations in this gene are associated with normosmic hypogonadotropic hypogonadism. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Feb 2016],

Western blot analysis of lysates from HuvEc cells, primary antibody was diluted at 1:1000, 4° over night





Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

