

## STX8 rabbit pAb

Cat No.: ES12886

For research use only

## Overview

Product Name STX8 rabbit pAb

Host species Rabbit

Applications WB;ELISA;IHC Species Cross-Reactivity Human; Mouse;Rat

**Recommended dilutions** WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 **Immunogen** Synthesized peptide derived from human STX8 AA

range: 42-92

**Specificity** This antibody detects endogenous levels of STX8 at

Human/Mouse/Rat

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name STX8
Gene Name STX8

Cellular localization Membrane; Single-pass type IV membrane protein.

Preferentially associated with the early endosome. To a lesser extent, also present in late endosome,

the plasma membrane and coated pits (By

similarity). .

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

**Observed band** 

Human Gene ID 9482 Human Swiss-Prot Number Q9UNK0

**Alternative Names** 

**Background** The gene is a member of the syntaxin family. The

encoded protein is involved in protein trafficking from early to late endosomes via vesicle fusion and

exocytosis. A related pseudogene has been


identified on chromosome 12. Alternative splicing





1 kDa 180 --140 --100 --75 --60 --45 --35 --25 --STX8 results in multiple transcript variants. [provided by RefSeq, May 2010],

Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

