

## RFX6 rabbit pAb

Cat No.: ES13408

For research use only

## Overview

Product Name RFX6 rabbit pAb

Host species Rabbit

Applications WB;ELISA;IHC Species Cross-Reactivity Human; Mouse

**Recommended dilutions** WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 **Immunogen** Synthesized peptide derived from human RFX6 AA

range: 264-314

**Specificity** This antibody detects endogenous levels of RFX6 at

Human/Mouse

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name RFX6

Gene Name RFX6 RFXDC1
Cellular localization Nucleus .

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

**Observed band** 

Human Gene ID 222546 Human Swiss-Prot Number Q8HWS3

**Alternative Names** 

**Background** The nuclear protein encoded by this gene is a

member of the regulatory factor X (RFX) family of transcription factors. Studies in mice suggest that

this gene is specifically required for the

differentiation of islet cells for the production of insulin, but not for the differentiation of pancreatic

polypeptide-producing cells. It regulates the

transcription factors involved in beta-cell maturation and function, thus, restricting the expression of the




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




beta-cell differentiation and specification genes. Mutations in this gene are associated with Mitchell-Riley syndrome, which is characterized by neonatal diabetes with pancreatic hypoplasia, duodenal and jejunal atresia, and gall bladder agenesis.[provided by RefSeq, Sep 2010],

Western blot analysis of lysates from SH-SY5Y cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950

