



# MORN2 rabbit pAb

Cat No.:ES14885

For research use only


## Overview

<b>Product Name</b>	MORN2 rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	IHC;IF
<b>Species Cross-Reactivity</b>	Human; Mouse
<b>Recommended dilutions</b>	IHC-p 1: 50-200
<b>Immunogen</b>	Synthesized peptide derived from human MORN2 AA range: 25-75
<b>Specificity</b>	This antibody detects endogenous levels of MORN2 at Human/Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	MORN2
<b>Gene Name</b>	MORN2 MOPT
<b>Cellular localization</b>	Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis. .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	
<b>Human Gene ID</b>	729967
<b>Human Swiss-Prot Number</b>	Q502X0
<b>Alternative Names</b>	
<b>Background</b>	





**ELK Biotechnology**



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950

[ELKbio@ELKbiotech.com](mailto:ELKbio@ELKbiotech.com)

[www.elkbiotech.com](http://www.elkbiotech.com)

23-2, No.388 Gaoxin 2nd Road,Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C