

PFK-2 car (phospho Ser483) rabbit pAb

Cat No.: ES1490

For research use only

Overview

Product Name PFK-2 car (phospho Ser483) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA **Species Cross-Reactivity** Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunohistochemistry:

1/100 - 1/300. ELISA: 1/40000. Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human PFKFB2 around the phosphorylation site of Ser483. AA range:451-500

Specificity Phospho-PFK-2 car (S483) Polyclonal Antibody detects

endogenous levels of PFK-2 car protein only when

phosphorylated at S483.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase

2

Gene Name PFKFB2
Cellular localization cytosol,

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 58kD
Human Gene ID 5208
Human Swiss-Prot Number 060825

Alternative Names PFKFB2; 6-phosphofructo-2-kinase/fructose-2;

6-bisphosphatase 2; 6PF-2-K/Fru-2,6-P2ase 2; PFK/FBPase 2; 6PF-2-K/Fru-2,6-P2ase heart-type

isozyme


Background The protein encoded by this gene is involved in both




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com



the synthesis and degradation of fructose-2,6-bisphosphate, a regulatory molecule that controls glycolysis in eukaryotes. The encoded protein has a 6-phosphofructo-2-kinase activity that catalyzes the synthesis of fructose-2,6-bisphosphate, and a fructose-2,6-biphosphatase activity that catalyzes the degradation of fructose-2,6-bisphosphate. This protein regulates fructose-2,6-bisphosphate levels in the heart, while a related enzyme encoded by a different gene regulates fructose-2,6-bisphosphate levels in the liver and muscle. This enzyme functions as a homodimer. Two transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],



Western Blot analysis of various cells using Phospho-PFK-2 car (S483) Polyclonal Antibody diluted at 1:1000




+86-27-59760950


Western Blot analysis of Hela cells using Phospho-PFK-2 car (S483) Polyclonal Antibody diluted at 1:1000







Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PFKFB2 (Phospho-Ser483) Antibody



+86-27-59760950

Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using PFKFB2 (Phospho-Ser483) Antibody. The picture on the right is blocked with the phospho peptide.

