



# IQCE rabbit pAb


Cat No.:ES15437

For research use only


## Overview

<b>Product Name</b>	IQCE rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;IHC
<b>Species Cross-Reactivity</b>	Human; Mouse
<b>Recommended dilutions</b>	WB 1:500-2000;IHC-p 1:50-300
<b>Immunogen</b>	Synthesized peptide derived from human IQCE AA range: 50-100
<b>Specificity</b>	This antibody detects endogenous levels of IQCE at Human/Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	IQCE
<b>Gene Name</b>	IQCE KIAA1023
<b>Cellular localization</b>	Cell projection, cilium membrane ; Peripheral membrane protein ; Cytoplasmic side . The EvC complex localizes at the base of cilia in the EvC zone of primary cilia in a EFCAB7-dependent manner. .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	
<b>Human Gene ID</b>	23288
<b>Human Swiss-Prot Number</b>	Q6IPM2
<b>Alternative Names</b>	
<b>Background</b>	





Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

