

ABCC12 rabbit pAb

Cat No.:ES1565

For research use only

Overview

Product Name ABCC12 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human MRP9. AA

range:691-740

Specificity ABCC12 Polyclonal Antibody detects endogenous

levels of ABCC12 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Multidrug resistance-associated protein 9

Gene Name ABCC12

Cellular localization Endoplasmic reticulum membrane ; Multi-pass

membrane protein.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band170kDHuman Gene ID94160Human Swiss-Prot NumberQ96J65

Alternative Names ABCC12; MRP9; Multidrug resistance-associated

protein 9; ATP-binding cassette sub-family C

member 12

Background This gene is a member of the superfamily of

ATP-binding cassette (ABC) transporters and the encoded protein contains two ATP-binding domains


and 12 transmembrane regions. ABC proteins




+86-27-59760950 ELKbio@ELKbiotech.com



transport various molecules across extra- and intracellular membranes. ABC genes are divided into seven distinct subfamilies: ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, and White. This gene is a member of the MRP subfamily which is involved in multi-drug resistance. This gene and another subfamily member are arranged head-to-tail on chromosome 16q12.1. Increased expression of this gene is associated with breast cancer. [provided by RefSeq, Jul 2008],



Western Blot analysis of various cells using ABCC12 Polyclonal Antibody diluted at 1:500



+86-27-59760950

Western blot analysis of lysates from 293, HepG2, and HUVEC cells, using MRP9 Antibody. The lane on the right is blocked with the synthesized peptide.

