

# FBX18 rabbit pAb

Cat No.:ES16506

For research use only


## Overview

Product Name	FBX18 rabbit pAb
Host species	Rabbit
Applications	WB
Species Cross-Reactivity	Human; Mouse
Recommended dilutions	WB 1: 500-2000
Immunogen	Synthesized peptide derived from human FBX18 AA range: 799-849
Specificity	This antibody detects endogenous levels of FBX18 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	FBX18
Gene Name	FBXO18 FBH1 FBX18
Cellular localization	Nucleus . Chromosome . Accumulates at sites of DNA damage or replication stress (PubMed:19736316, PubMed:23677613). PCNA is required for localization to DNA damage sites (PubMed:23677613). Localizes to the nucleoplasm in absence of DNA damage (PubMed:2367
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	
Human Gene ID	84893
Human Swiss-Prot Number	Q8NFZ0
Alternative Names	
Background	This gene encodes a member of the F-box protein family, members of which are characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of





ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into three classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbx class. It contains an F-box motif and seven conserved helicase motifs, and has both DNA-dependent ATPase and DNA unwinding activities. Alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Jul 2008],



Western blot analysis of lysates from HuvEc cells, primary antibody was diluted at 1:1000, 4° over night

