

CYH4 rabbit pAb

Cat No.: ES17132

For research use only

Overview

Product Name CYH4 rabbit pAb

Host species Rabbit
Applications WB

Species Cross-Reactivity Human; Mouse Recommended dilutions WB 1: 500-2000

Immunogen Synthesized peptide derived from human CYH4 AA

range: 150-200

Specificity This antibody detects endogenous levels of CYH4 at

Human/Mouse

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name CYH4

Gene Name CYTH4 CYT4 PSCD4

Cellular localizationCell membrane ; Peripheral membrane protein .PurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

Observed band

Human Gene ID 27128 Human Swiss-Prot Number Q9UIA0

Alternative Names

Background This gene encodes a member of the PSCD family of

proteins, which have an N-terminal coiled-coil motif, a central Sec7 domain, and a C-terminal pleckstrin homology (PH) domain. The coiled-coil motif is involved in homodimerization, the Sec7 domain contains guanine-nucleotide exchange protein (GEP)

activity, and the PH domain interacts with

phospholipids and is responsible for association of PSCDs with membranes. Members of this family



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



1 kDa
180 -140 -140 -100 -75 -60 -45 -25 -15 -10 --

function as GEPs for ADP-ribosylation factors (ARFs), which are guanine nucleotide-binding proteins involved in vesicular trafficking pathways. This protein exhibits GEP activity in vitro with ARF1 and ARF5, but is inactive with ARF6. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Dec 2015],

Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night

