

## **CBG** rabbit pAb

## Cat No.:ES1875

For research use only

## Overview

Product Name	CBG rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not
	yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human GBA3. AA
	range:291-340
Specificity	CBG Polyclonal Antibody detects endogenous levels
	of CBG protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Cytosolic beta-glucosidase
Gene Name	GBA3
Cellular localization	Cytoplasm, cytosol .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	54-58kD
Human Gene ID	57733
Human Swiss-Prot Number	Q9H227
Alternative Names	GBA3; CBG; CBGL1; Cytosolic beta-glucosidase;
	Cytosolic beta-glucosidase-like protein 1
Background	The protein encoded by this gene is an enzyme that
	can hydrolyze several types of glycosides. This gene
	is a polymorphic pseudogene, with the most
	common allele being the functional allele that
	encodes the full-length protein. Some individuals, as
	represented by the reference genome allele, contain



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



HuvEc

(kD)

170-

130-

95-

72-

55-

GBA3--=

(kD)

117-

85-

48-

34-

26-

19-

HuvEC HuvEC

-- 117

-- 85

-- 48

-- 34

-- 26 -- 19 (kD)

GBA3

a single nucleotide polymorphism that results in a premature stop codon in the coding region, and therefore this allele is pseudogenic due to the failure to produce a functional full-length protein. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Mar 2013],

Western Blot analysis of various cells using CBG Polyclonal Antibody diluted at 1:500

Western blot analysis of lysates from HUVEC cells, using GBA3 Antibody. The lane on the right is blocked with the synthesized peptide.

Western blot analysis of the lysates from HeLa cells using GBA3 antibody.



www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C