

Histone H2A (Acetyl Lys6) rabbit pAb

Cat No.:ES20072

For research use only

Overview

| Product Name | Histone H2A (Acetyl Lys6) rabbit pAb |
|--------------------------|--|
| Host species | Rabbit |
| Applications | WB; ELISA |
| Species Cross-Reactivity | Human;Mouse;Rat |
| Recommended dilutions | WB 1:1000-2000 ELISA 1:5000-20000 |
| Immunogen | Synthesized peptide derived from human Histone |
| | H2A (Acetyl Lys6) |
| Specificity | This antibody detects endogenous levels of |
| | Human,Mouse,Rat Histone H2A (Acetyl Lys6) |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20 $^\circ\!{ m C}$. Avoid repeated freeze-thaw cycles. |
| Protein Name | Histone H2A (Acetyl Lys6) |
| Gene Name | HIST1H2AG H2AFP; HIST1H2AI H2AFC; HIST1H2AK |
| | H2AFD; HIST1H2AL H2AFI; HIST1H2AM H2AFN |
| Cellular localization | Nucleus. Chromosome. |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 15kD |
| Human Gene ID | 8329 |
| Human Swiss-Prot Number | P0C0S8 |
| Alternative Names | Histone H2A type 1 (H2A.1;Histone H2A/p) |
| Background | Histones are basic nuclear proteins that are |
| | responsible for the nucleosome structure of the |
| | chromosomal fiber in eukaryotes. Two molecules of |
| | each of the four core histones (H2A, H2B, H3, and |
| | H4) form an octamer, around which approximately |
| | 146 bp of DNA is wrapped in repeating units, called |
| | nucleosomes. The linker histone, H1, interacts with |
| | linker DNA between nucleosomes and functions in |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015],



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C