

## COL5A1 rabbit pAb

## Cat No.:ES2025

For research use only

## Overview

Product Name	COL5A1 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300.
	Immunofluorescence: 1/200 - 1/1000. ELISA:
	1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human Collagen V alpha1. AA
	range:301-350
Specificity	COL5A1 Polyclonal Antibody detects endogenous
	levels of COL5A1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Collagen alpha-1(V) chain
Gene Name	COL5A1
Cellular localization	Secreted, extracellular space, extracellular matrix .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	200kD
Human Gene ID	1289
Human Swiss-Prot Number	P20908
Alternative Names	COL5A1; Collagen alpha-1(V) chain
Background	This gene encodes an alpha chain for one of the low
	abundance fibrillar collagens. Fibrillar collagen
	molecules are trimers that can be composed of one
	or more types of alpha chains. Type V collagen is
	found in tissues containing type I collagen and



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C




appears to regulate the assembly of heterotypic fibers composed of both type I and type V collagen. This gene product is closely related to type XI collagen and it is possible that the collagen chains of types V and XI constitute a single collagen type with tissue-specific chain combinations. The encoded procollagen protein occurs commonly as the heterotrimer

pro-alpha1(V)-pro-alpha1(V)-pro-alpha2(V). Mutations in this gene are associated with Ehlers-Danlos syndrome, types I and II. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, May 2013],

Western Blot analysis of various cells using COL5A1 Polyclonal Antibody



Immunofluorescence analysis of HeLa cells, using Collagen V alpha1 Antibody. The picture on the right is blocked with the synthesized peptide.





+86-27-59760950


ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Immunohistochemistry analysis of paraffin-embedded human heart tissue, using Collagen V alpha1 Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from HeLa, and HUVEC, and HepG2 cells, using Collagen V alpha1 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C