

## COP ζ1 rabbit pAb

## Cat No.:ES2030

For research use only

## Overview

Product Name	COP ζ1 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Monkey
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human COPZ1. AA range:11-60
Specificity	COP ζ1 Polyclonal Antibody detects endogenous
	levels of COP ζ1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Coatomer subunit zeta-1
Gene Name	COPZ1
Cellular localization	Cytoplasm . Golgi apparatus membrane ; Peripheral
	membrane protein ; Cytoplasmic side . Cytoplasmic
	vesicle, COPI-coated vesicle membrane ; Peripheral
	membrane protein ; Cytoplasmic side . The
	coatomer is cytoplasmic or polymerized on the
	cytoplasmic side
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	20kD
Human Gene ID	22818
Human Swiss-Prot Number	P61923
Alternative Names	COPZ1; COPZ; CGI-120; HSPC181; Coatomer subunit
	zeta-1; Zeta-1-coat protein; Zeta-1 COP
Background	This gene encodes a subunit of the cytoplasmic



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



coatamer protein complex, which is involved in autophagy and intracellular protein trafficking. The coatomer protein complex is comprised of seven subunits and functions as the coat protein of coat protein complex (COP)I-vesicles. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2012],

Western Blot analysis of various cells using COP  $\zeta$ 1 Polyclonal Antibody diluted at 1:500

Immunohistochemistry analysis of paraffin-embedded human cervix carcinoma tissue, using COPZ1 Antibody. The picture on the right is blocked with the synthesized peptide.



Jurka

(kD)

117-85-

48-

34-

26-

19-

Western blot analysis of lysates from Jurkat and COS cells, using COPZ1 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C