

CPE rabbit pAb

Cat No.:ES2039

For research use only

Overview

Product Name CPE rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human CPE. AA range:271-320

Specificity CPE Polyclonal Antibody detects endogenous levels

of CPE protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Carboxypeptidase E

Gene Name CPE

Cellular localization [Isoform 1]: Cytoplasmic vesicle, secretory vesicle.

Cytoplasmic vesicle, secretory vesicle membrane; Peripheral membrane protein . Secreted . Associated with the secretory granule membrane through direct binding to lipid rafts in intragranular

conditions. .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 53kD
Human Gene ID 1363
Human Swiss-Prot Number P16870

Alternative Names CPE; Carboxypeptidase E; CPE; Carboxypeptidase H;

CPH; Enkephalin convertase;

Prohormone-processing carboxypeptidase


Background carboxypeptidase E(CPE) Homo sapiens This



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com




gene encodes a member of the M14 family of metallocarboxypeptidases. The encoded preproprotein is proteolytically processed to generate the mature peptidase. This peripheral membrane protein cleaves C-terminal amino acid residues and is involved in the biosynthesis of peptide hormones and neurotransmitters, including insulin. This protein may also function independently of its peptidase activity, as a neurotrophic factor that promotes neuronal survival, and as a sorting receptor that binds to regulated secretory pathway proteins, including prohormones. Mutations in this gene are implicated in type 2 diabetes. [provided by RefSeq, Nov 2015],



27

Western Blot analysis of various cells using CPE Polyclonal Antibody



+86-27-59760950


Western Blot analysis of K562 cells using CPE Polyclonal Antibody




ELKbio@ELKbiotech.com

www.elkbiotech.com





Western blot analysis of lysates from RAW264.7 and NIH/3T3 cells, using CPE Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HeLa cells using CPE antibody.

