

CTPS rabbit pAb

Cat No.: ES2070

For research use only

Overview

Product Name CTPS rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human CTPS. AA range:60-109

Specificity CTPS Polyclonal Antibody detects endogenous levels

of CTPS protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name CTP synthase 1

Gene Name CTPS1

Cellular localization Cytoplasm, cytosol . Mainly cytosolic but when

active detected in long filamentous structures (PubMed:25223282). Co-localizes with TNK2 in the

cytosolic filaments (By similarity). .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 67kD
Human Gene ID 1503
Human Swiss-Prot Number P17812

Alternative Names CTPS1; CTPS; CTP synthase 1; CTP synthetase 1;

UTP--ammonia ligase 1

Background This gene encodes an enzyme responsible for the

catalytic conversion of UTP (uridine triphosphate) to


CTP (cytidine triphospate). This reaction is an





important step in the biosynthesis of phospholipids and nucleic acids. Activity of this proten is important in the immune system, and loss of function of this gene has been associated with immunodeficiency. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2014],

Western Blot analysis of various cells using CTPS Polyclonal Antibody diluted at 1:500




95-

72-

55-

Western Blot analysis of HuvEc cells using CTPS Polyclonal Antibody diluted at 1:500



Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbe





Immunohistochemistry analysis of CTPS antibody in paraffin-embedded human liver carcinoma tissue.

