

CYP2J2 rabbit pAb

Cat No.:ES2104

For research use only

Overview

Product Name	CYP2J2 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Monkey
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300.
	Immunofluorescence: 1/200 - 1/1000. ELISA:
	1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human Cytochrome P450 2J2.
	AA range:231-280
Specificity	CYP2J2 Polyclonal Antibody detects endogenous
	levels of CYP2J2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Cytochrome P450 2J2
Gene Name	CYP2J2
Cellular localization	Endoplasmic reticulum membrane; Peripheral
	membrane protein. Microsome membrane ;
	Peripheral membrane protein.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	58kD
Human Gene ID	1573
Human Swiss-Prot Number	P51589
Alternative Names	CYP2J2; Cytochrome P450 2J2; Arachidonic acid
	epoxygenase; CYPIIJ2
Background	This gene encodes a member of the cytochrome
	P450 superfamily of enzymes. The cytochrome P450



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is thought to be the predominant enzyme responsible for epoxidation of endogenous arachidonic acid in cardiac tissue. Multiple transcript variants have been found for this gene. [provided by RefSeq, Jan 2016],

Western Blot analysis of various cells using CYP2J2 Polyclonal Antibody

Immunofluorescence analysis of HepG2 cells, using Cytochrome P450 2J2 Antibody. The picture on the right is blocked with the synthesized peptide.



HT29

(kD)

170-


130-

95-

72-

55-

Immunohistochemistry analysis of paraffin-embedded human heart tissue, using Cytochrome P450 2J2 Antibody. The picture on the right is blocked with the synthesized peptide.




+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Western blot analysis of lysates from A549 and COS7 cells, using Cytochrome P450 2J2 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C