

ERCC4 rabbit pAb

Cat No.: ES2292

For research use only

Overview

Product Name ERCC4 rabbit pAb

Host species Rabbit WB;ELISA **Applications Species Cross-Reactivity** Human; Mouse

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

The antiserum was produced against synthesized **Immunogen**

peptide derived from human XPF. AA range:801-850

Specificity ERCC4 Polyclonal Antibody detects endogenous

levels of ERCC4 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage**

Protein Name DNA repair endonuclease XPF

Gene Name ERCC4

Cellular localization Nucleus . Chromosome . Localizes to sites of DNA

damage...

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 103kD **Human Gene ID** 2072 **Human Swiss-Prot Number** Q92889

Alternative Names ERCC4; ERCC11; XPF; DNA repair endonuclease XPF;

> DNA excision repair protein ERCC-4; DNA repair protein complementing XP-F cells; Xeroderma pigmentosum group F-complementing protein The protein encoded by this gene forms a complex


Background

with ERCC1 and is involved in the 5' incision made during nucleotide excision repair. This complex is a structure specific DNA repair endonuclease that



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com







26-

interacts with EME1. Defects in this gene are a cause of xeroderma pigmentosum complementation group F (XP-F), or xeroderma pigmentosum VI (XP6).[provided by RefSeq, Mar 2009],

Western Blot analysis of various cells using ERCC4
Polyclonal Antibody cells nucleus extracted by Minute TM
Cytoplasmic and Nuclear Fractionation kit
(SC-003,Inventbiotech,MN,USA).



Western Blot analysis of 293 cells using ERCC4 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Western blot analysis of lysates from 293 cells, using XPF Antibody. The lane on the right is blocked with the synthesized peptide.

