

Frizzled-1 rabbit pAb

Cat No.:ES2371

For research use only

Overview

| Product Name | Frizzled-1 rabbit pAb |
|--------------------------|---|
| Host species | Rabbit |
| Applications | WB;IF;ELISA |
| Species Cross-Reactivity | Human;Rat;Mouse;;Rice |
| Recommended dilutions | Western Blot: 1/500 - 1/2000. Immunofluorescence: |
| | 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in |
| | other applications. |
| Immunogen | The antiserum was produced against synthesized |
| | peptide derived from human FZD1. AA range:15-64 |
| Specificity | Frizzled-1 Polyclonal Antibody detects endogenous |
| | levels of Frizzled-1 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | Frizzled-1 |
| Gene Name | FZD1 |
| Cellular localization | Cell membrane ; Multi-pass membrane protein . |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 71kD |
| Human Gene ID | 8321 |
| Human Swiss-Prot Number | Q9UP38 |
| Alternative Names | FZD1; Frizzled-1; Fz-1; hFz1; FzE1 |
| Background | frizzled class receptor 1(FZD1) Homo sapiens |
| | Members of the 'frizzled' gene family encode |
| | 7-transmembrane domain proteins that are |
| | receptors for Wnt signaling proteins. The FZD1 |
| | protein contains a signal peptide, a cysteine-rich |
| | domain in the N-terminal extracellular region, 7 |
| | transmembrane domains, and a C-terminal PDZ |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



domain-binding motif. The FZD1 transcript is expressed in various tissues. [provided by RefSeq, Jul 2008],

Western Blot analysis of various cells using Frizzled-1 Polyclonal Antibody

Western blot analysis of lysates from HeLa and COLO205 cells, using FZD1 Antibody. The lane on the right is blocked with the synthesized peptide.



HeLa

(kD)

117-85-

48-

34-

26-

19-

FZD1-

HeLa COLO COLO

-- 117

-- 85

-- 48 -- 34

-- 26

-- 19

Western blot analysis of the lysates from HT-29 cells using FZD1 antibody.



www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C