

## hnRNP G rabbit pAb

## Cat No.:ES2549

For research use only

## **Overview**

Product Name	hnRNP G rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300.
	Immunofluorescence: 1/200 - 1/1000. ELISA:
	1/40000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human hnRNP G. AA
	range:6-55
Specificity	hnRNP G Polyclonal Antibody detects endogenous
	levels of hnRNP G protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	RNA-binding motif protein X chromosome
Gene Name	RBMX
Cellular localization	Nucleus . Component of ribonucleosomes. Localizes
	in numerous small granules in the nucleus.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	40kD
Human Gene ID	27316
Human Swiss-Prot Number	
Alternative Names	RBMX; HNRPG; RBMXP1; RNA-binding motif
	protein; X chromosome; Glycoprotein p43;
	Heterogeneous nuclear ribonucleoprotein G; hnRNP
	G
Background	This gene belongs to the RBMY gene family which
+86-27-59760950	ELKbio@ELKbiotech.com www.elkbiotech.com




23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C




includes candidate Y chromosome spermatogenesis genes. This gene, an active X chromosome homolog of the Y chromosome RBMY gene, is widely expressed whereas the RBMY gene evolved a male-specific function in spermatogenesis. Pseudogenes of this gene, found on chromosomes 1, 4, 9, 11, and 6, were likely derived by retrotransposition from the original gene. Alternatively spliced transcript variants encoding different isoforms have been identified. A snoRNA gene (SNORD61) is found in one of its introns. [provided by RefSeq, Sep 2009],

Western Blot analysis of various cells using hnRNP G Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Immunofluorescence analysis of HepG2 cells, using hnRNP G Antibody. The picture on the right is blocked with the synthesized peptide.






+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C





Western blot analysis of lysates from HeLa cells, using hnRNP G Antibody. The lane on the right is blocked with the synthesized peptide.

Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200(4° overnight.3,Secondary antibody was diluted at 1:200(room temperature, 45min).





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C