

hnRNP L rabbit pAb

Cat No.: ES2552

For research use only

Overview

Product Name hnRNP L rabbit pAb

Host species Rabbit

ApplicationsWB;IHC;IF;ELISASpecies Cross-ReactivityHuman;Mouse

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. The antiserum was produced against synthesized

Immunogen The antiserum was produced against synthesized

peptide derived from human hnRNP L. AA

range:61-110

Specificity hnRNP L Polyclonal Antibody detects endogenous

levels of hnRNP L protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Heterogeneous nuclear ribonucleoprotein L

Gene Name HNRNPL

Cellular localization Nucleus, nucleoplasm . Cytoplasm . Localized in

cytoplasmic mRNP granules containing untranslated mRNAs. These granules are not identical with P

bodies or stress granules. .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 60kD
Human Gene ID 3191
Human Swiss-Prot Number P14866

Alternative Names HNRNPL; HNRPL; P/OKcl.14; Heterogeneous nuclear

ribonucleoprotein L; hnRNP L


Background Heterogeneous nuclear RNAs (hnRNAs) which




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com



include mRNA precursors and mature mRNAs are associated with specific proteins to form heterogenous ribonucleoprotein (hnRNP) complexes. Heterogeneous nuclear ribonucleoprotein L is among the proteins that are stably associated with hnRNP complexes and along with other hnRNP proteins is likely to play a major role in the formation, packaging, processing, and function of mRNA. Heterogeneous nuclear ribonucleoprotein L is present in the nucleoplasm as part of the HNRP complex. HNRP proteins have also been identified outside of the nucleoplasm. Exchange of hnRNP for mRNA-binding proteins accompanies transport of mRNA from the nucleus to the cytoplasm. Since HNRP proteins have been shown to shuttle between the nucleus and the cytoplasm, it is possible that they also have cytoplasmic functions. Two transcript variants encoding different is



Western Blot analysis of various cells using hnRNP L Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Western Blot analysis of HuvEc cells using hnRNP L Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).





+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com







Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using hnRNP L Antibody. The picture on the right is blocked with the synthesized peptide.



+86-27-59760950

