

IL-13Rα1 rabbit pAb

Cat No.:ES2608

For research use only

Overview

Immunogen

Product Name IL-13Rα1 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA:

1/10000. Not yet tested in other applications.
The antiserum was produced against synthesized

peptide derived from human IL-13R/CD213 alpha1.

AA range:371-420

Specificity IL-13Rα1 Polyclonal Antibody detects endogenous

levels of IL-13Rα1 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Interleukin-13 receptor subunit alpha-1

Gene Name IL13RA1

Cellular localizationMembrane; Single-pass type I membrane protein.PurificationThe antibody was affinity-purified from rabbit
antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 49kD
Human Gene ID 3597
Human Swiss-Prot Number P78552

Alternative Names IL13RA1; IL13R; IL13RA; Interleukin-13 receptor

subunit alpha-1; IL-13 receptor subunit alpha-1; IL-13R subunit alpha-1; IL-13R-alpha-1; IL-13RA1; Cancer/testis antigen 19; CT19; CD antigen CD213a1 The protein encoded by this gene is a subunit of the

Background The protein encoded by this gene is a subunit of

interleukin 13 receptor. This subunit forms a




+86-27-59760950 ELKbio@ELKbiotech.com




receptor complex with IL4 receptor alpha, a subunit shared by IL13 and IL4 receptors. This subunit serves as a primary IL13-binding subunit of the IL13 receptor, and may also be a component of IL4 receptors. This protein has been shown to bind tyrosine kinase TYK2, and thus may mediate the signaling processes that lead to the activation of JAK1, STAT3 and STAT6 induced by IL13 and IL4. [provided by RefSeq, Jul 2008],


Western Blot analysis of various cells using IL-13R α 1 Polyclonal Antibody diluted at 1:1000



Immunofluorescence analysis of NIH/3T3 cells, using IL-13R/CD213 alpha1 Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using IL-13R/CD213 alpha1 Antibody. The picture on the right is blocked with the synthesized peptide.



+86-27-59760950







Western blot analysis of lysates from COLO, HT-29, A549, and 293 cells, using IL-13R/CD213 alpha1 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950