

MSY2 rabbit pAb

Cat No.: ES2862

For research use only

Overview

Product Name MSY2 rabbit pAb

Host species Rabbit

ApplicationsWB;IHC;IF;ELISASpecies Cross-ReactivityHuman;Mouse

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. The antiserum was produced against synthesized

Immunogen The antiserum was produced against synthesized

peptide derived from human YBOX2. AA

range:281-330

Specificity MSY2 Polyclonal Antibody detects endogenous

levels of MSY2 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Y-box-binding protein 2

Gene Name YBX2

Cellular localization Cytoplasm . Nucleus .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 38kD
Human Gene ID 51087
Human Swiss-Prot Number Q9Y2T7

Alternative Names YBX2; CSDA3; MSY2; Y-box-binding protein 2;

Contrin; DNA-binding protein C; Dbpc; Germ

cell-specific Y-box-binding protein; MSY2 homolog

Background This gene encodes a nucleic acid binding protein

which is highly expressed in germ cells. The encoded protein binds to a Y-box element in the promoters of




+86-27-59760950 ELKbio@ELKbiotech.com



certain genes but also binds to mRNA transcribed from these genes. Pseudogenes for this gene are located on chromosome 10 and 15. [provided by RefSeq, Feb 2012],

Western Blot analysis of various cells using MSY2 Polyclonal Antibody



-- 117

(kD)

Western blot analysis of lysates from COLO205 cells, using YBOX2 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

