

Neurotrimin rabbit pAb

Cat No.: ES2935

For research use only

Overview

Product Name Neurotrimin rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. The antiserum was produced against synthesized

Immunogen The antiserum was produced against synthesized

peptide derived from human NT. AA range:235-284

Specificity Neurotrimin Polyclonal Antibody detects

endogenous levels of Neurotrimin protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Neurotrimin

Gene Name NTM

Cellular localizationCell membrane; Lipid-anchor, GPI-anchor.PurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 42kD
Human Gene ID 50863
Human Swiss-Prot Number Q9P121

Alternative Names NTM; IGLON2; NT; Neurotrimin; hNT; IgLON family

member 2

Background This gene encodes a member of the IgLON (LAMP,

OBCAM, Ntm) family of immunoglobulin (Ig) domain-containing glycosylphosphatidylinositol (GPI)-anchored cell adhesion molecules. The encoded protein may promote neurite outgrowth and adhesion via a homophilic mechanism. This




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




gene is closely linked to a related family member, opioid binding protein/cell adhesion molecule-like (OPCML), on chromosome 11. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2009],

Western Blot analysis of various cells using Neurotrimin Polyclonal Antibody



Western blot analysis of lysates from HUVEC, HepG2, and Jurkat cells, using NT Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using NT antibody.

