

## OX2R rabbit pAb

## Cat No.:ES3089

For research use only

## Overview

Product Name	OX2R rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not
	yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
-	peptide derived from human MOX2R. AA
	range:241-290
Specificity	OX2R Polyclonal Antibody detects endogenous
	levels of OX2R protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Cell surface glycoprotein CD200 receptor 1
Gene Name	CD200R1
Cellular localization	[Isoform 1]: Cell membrane; Single-pass type I
	membrane protein.; [Isoform 4]: Cell membrane;
	Single-pass type I membrane protein.; [Isoform 2]:
	Secreted.; [Isoform 3]: Secreted.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	37kD
Human Gene ID	131450
Human Swiss-Prot Number	Q8TD46
Alternative Names	CD200R1; CD200R; CRTR2; MOX2R; OX2R; Cell
	surface glycoprotein CD200 receptor 1; CD200 cell
	surface glycoprotein receptor; Cell surface
	glycoprotein OX2 receptor 1
Background	This gene encodes a receptor for the OX-2
7	



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



KDa

120

90

50

39

27

(kD)

117-85-

48-

34-

26-


19-

OX2R

HepG2

membrane glycoprotein. Both the receptor and substrate are cell surface glycoproteins containing two immunoglobulin-like domains. This receptor is restricted to the surfaces of myeloid lineage cells and the receptor-substrate interaction may function as a myeloid downregulatory signal. Mouse studies of a related gene suggest that this interaction may control myeloid function in a tissue-specific manner. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jul 2008],

Western Blot analysis of K562 cells using OX2R Polyclonal Antibody






ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Western blot analysis of lysates from K562 cells, using MOX2R Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C