

PFK-2 liv rabbit pAb

Cat No.:ES3183

For research use only

Overview

Product Name	PFK-2 liv rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species	Human;Mouse;Rat
Cross-Reactivity	
Recommended	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100
dilutions	- 1/300. ELISA: 1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide
	derived from human PFKFB1. AA range:406-455
Specificity	PFK-2 liv Polyclonal Antibody detects endogenous levels of
	PFK-2 liv protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02%
	sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase 1
Gene Name	PFKFB1
Cellular localization	cytosol,6-phosphofructo-2-kinase/fructose-2,6-biphosphatase
	complex,
Purification	The antibody was affinity-purified from rabbit antiserum by
	affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	58kD
Human Gene ID	5207
Human Swiss-Prot	P16118
Number	
Alternative Names	PFKFB1; F6PK; PFRX; 6-phosphofructo-2-kinase/fructose-2;
	6-bisphosphatase 1; 6PF-2-K/Fru-2,6-P2ase 1; PFK/FBPase 1;
	6PF-2-K/Fru-2,6-P2ase liver isozyme
Background	This gene encodes a member of the family of bifunctional
	6-phosphofructo-2-kinase:fructose-2,6-biphosphatase
	enzymes. The enzyme forms a homodimer that catalyzes both
	the synthesis and degradation of fructose-2,6-biphosphate



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



using independent catalytic domains.

Fructose-2,6-biphosphate is an activator of the glycolysis pathway and an inhibitor of the gluconeogenesis pathway. Consequently, regulating fructose-2,6-biphosphate levels through the activity of this enzyme is thought to regulate glucose homeostasis. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Nov 2012],






+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C





Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C