

## β-1,3-Gal-T1 rabbit pAb

## Cat No.:ES3747

For research use only

## Overview

Product Name Host species Applications Species Cross-Reactivit y	β-1,3-Gal-T1 rabbit pAb Rabbit WB;ELISA Human;Mouse
, Recommende	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in
d dilutions	other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human B3GALT1. AA range:61-110
Specificity	β-1,3-Gal-T1 Polyclonal Antibody detects endogenous levels of β-1,3-Gal-T1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Beta-1,3-galactosyltransferase 1
Gene Name	B3GALT1
Cellular	Golgi apparatus membrane ; Single-pass type II membrane protein .
localization	
Purification	The antibody was affinity-purified from rabbit antiserum by
	affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed	36kD
band	
Human Gene ID	8708
Human	Q9Y5Z6
Swiss-Prot	
Number	
Alternative Names	B3GALT1; Beta-1; 3-galactosyltransferase 1; Beta-1,3-GalTase 1; Beta3Gal-T1; Beta3GalT1; UDP-galactose:beta-N-acetyl-glucosamine-beta-1,3-galactosyltransf
	Senerce in accel. Braccontine were the Balaccoshirano



+86-27-59760950 ELKbio@ELKbiotech.com

om www

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



## erase 1

Background

This gene is a member of the beta-1,3-galactosyltransferase (beta3GalT) gene family. This family encodes type II membrane-bound glycoproteins with diverse enzymatic functions using different donor substrates (UDP-galactose and UDP-N-acetylglucosamine) and different acceptor sugars (N-acetylglucosamine, galactose, N-acetylgalactosamine). The beta3GalT genes are distantly related to the Drosophila Brainiac gene and have the protein coding sequence contained in a single exon. The beta3GalT proteins also contain conserved sequences not found in the beta4GalT or alpha3GalT proteins. The carbohydrate chains synthesized by these enzymes are designated as type 1, whereas beta4GalT enzymes synthesize type 2 carbohydrate chains. The ratio of type 1:type 2 chains changes during embryogenesis. By sequence similarity, the beta3GalT genes fall into at least two groups: beta3GalT4 and 4 other beta3



Western Blot analysis of various cells using  $\beta$ -1,3-Gal-T1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysates from HUVEC, COLO, Jurkat, and HepG2 cells, using B3GALT1 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Western blot analysis of the lysates from HUVECcells using B3GALT1 antibody.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C