

PAOX rabbit pAb

Cat No.:ES3828

For research use only


Overview

Product Name	PAOX rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/40000. Not yet tested in other applications.
Immunogen	Synthesized peptide derived from PAOX . at AA range: 260-340
Specificity	PAOX Polyclonal Antibody detects endogenous levels of PAOX protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Peroxisomal N(1)-acetyl-spermine/spermidine oxidase
Gene Name	PAOX
Cellular localization	Peroxisome . Cytoplasm .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	70kD
Human Gene ID	196743
Human Swiss-Prot Number	Q6QHF9
Alternative Names	PAOX; PAO; Peroxisomal N(1)-acetyl-spermine/spermidine oxidase; Polyamine oxidase
Background	catalytic activity:N(1),N(12)-diacetylspermine + O(2) + H(2)O = N(1)-acetylspermidine + 3-acetamidobutanal + H(2)O(2).,catalytic activity:N(1)-acetylspermidine + O(2) + H(2)O = putrescine + 3-acetamidopropanal +






H₂O₂.,catalytic activity:N(1)-acetylspermine + O₂ + H₂O = spermidine + 3-acetamidopropanal + H₂O₂.,cofactor:Binds 1 FAD per subunit.,function:Flavoenzyme which catalyzes the oxidation of N(1)-acetylspermine to spermidine and is thus involved in the polyamine back-conversion. Can also oxidize N(1)-acetylspermidine to putrescine. Substrate specificity: N(1)-acetylspermine = N(1)-acetylspermidine > N(1),N(12)-diacetylspermine >> spermine. Does not oxidize spermidine. Plays an important role in the regulation of polyamine intracellular concentration and has the potential to act as a determinant of cellular sensitivity to the antitumor polyamine analogs.,induction:By polyamine analogs.,miscellaneous:Oxidizes N(1)-acetylated polyamines on the exo-side of their N(4)-amino groups. Plant PAO oxidizes spermine on the endo-side of the N(4)-nitrogen.,pathway:Amine and polyamine metabolism; spermine metabolism.,similarity:Belongs to the flavin monoamine oxidase family.,subunit:Monomer.,tissue specificity:Widely expressed. Not detected in spleen. Expressed at lower level in neoplastic tissues.,



Western Blot analysis of extracts from rat stomach, K562 cells, using PAOX Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100

