

Actin-α cardiac muscle rabbit pAb

Cat No.: ES3840

For research use only

Overview

Product Name Actin-α cardiac muscle rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300

ELISA: 1/20000. Not yet tested in other applications.

Immunogen Synthesized peptide derived from Actin- α cardiac

muscle . at AA range: 1-80

Specificity Actin-α cardiac muscle Polyclonal Antibody detects

endogenous levels of Actin-α cardiac muscle

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Actin alpha cardiac muscle 1

Gene Name ACTC1

Cellular localization Cytoplasm, cytoskeleton.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 42kD
Human Gene ID 70
Human Swiss-Prot Number P68032

Alternative Names ACTC1; ACTC; Actin, alpha cardiac muscle 1;

Alpha-cardiac actin

Background Actins are highly conserved proteins that are

involved in various types of cell motility.

Polymerization of globular actin (G-actin) leads to a


structural filament (F-actin) in the form of a two-stranded helix. Each actin can bind to four others. The protein encoded by this gene belongs to




+86-27-59760950 ELKbio@ELKbiotech.com wv



the actin family which is comprised of three main groups of actin isoforms, alpha, beta, and gamma. The alpha actins are found in muscle tissues and are a major constituent of the contractile apparatus. Defects in this gene have been associated with idiopathic dilated cardiomyopathy (IDC) and familial hypertrophic cardiomyopathy (FHC). [provided by RefSeq, Jul 2008],



Western Blot analysis of mouse heart, mouse lung cells using Actin- α cardiac muscle Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded rat-muscle, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded rat-muscle, antibody was diluted at 1:100



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



Immunohistochemical analysis of paraffin-embedded mouse-muscle, antibody was diluted at 1:100



+86-27-59760950