

JAM-B rabbit pAb

Cat No.:ES3845

For research use only

Overview

Product Name JAM-B rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300

ELISA: 1/20000. Not yet tested in other applications.

Immunogen Synthesized peptide derived from the Internal

region of human JAM-B.

Specificity JAM-B Polyclonal Antibody detects endogenous

levels of JAM-B protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Junctional adhesion molecule B

Gene Name JAM2

Cellular localization Cell membrane ; Single-pass type I membrane

protein . Cell junction . Cell junction, tight junction . Localized at tight junctions of both epithelial and endothelial cells (By similarity). Specifically localized

within the somatodendritic compartment of

neurons and excluded from the axon (By similarity). .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 33kD
Human Gene ID 58494
Human Swiss-Prot Number P57087

Alternative Names JAM2; C21orf43; VEJAM; Junctional adhesion

molecule B; JAM-B; Junctional adhesion molecule 2; JAM-2; Vascular endothelial junction-associated

molecule; VE-JAM; CD322




+86-27-59760950 ELKbio@ELKbiotech.com




Background

This gene belongs to the immunoglobulin superfamily, and the junctional adhesion molecule (JAM) family. The protein encoded by this gene is a type I membrane protein that is localized at the tight junctions of both epithelial and endothelial cells. It acts as an adhesive ligand for interacting with a variety of immune cell types, and may play a role in lymphocyte homing to secondary lymphoid organs. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2012],

Western Blot analysis of K562 cells using JAM-B Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded rat-lung, antibody was diluted at 1:100










Immunohistochemical analysis of paraffin-embedded rat-lung, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-lung, antibody was diluted at 1:100

