



CD177 rabbit pAb

Cat No.:ES4013

For research use only

Overview


Product Name	CD177 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human CD177. AA range:361-410
Specificity	CD177 Polyclonal Antibody detects endogenous levels of CD177 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	CD177 antigen
Gene Name	CD177
Cellular localization	Cell membrane ; Lipid-anchor, GPI-anchor . Membrane raft ; Lipid-anchor, GPI-like-anchor . Secreted . Cytoplasmic granule membrane . Cell projection, lamellipodium . Cell surface expression on neutrophils is increased upon TNF-alpha, fMLP or CXCL8/IL8-med
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	46kD
Human Gene ID	57126
Human Swiss-Prot Number	Q8N6Q3
Alternative Names	CD177; NB1; PRV1; CD177 antigen; Human neutrophil alloantigen 2a; HNA-2a; NB1 glycoprotein; NB1 GP; Polycythemia rubra vera




Background

protein 1; PRV-1; CD177

This gene encodes a glycosyl-phosphatidylinositol (GPI)-linked cell surface glycoprotein that plays a role in neutrophil activation. The protein can bind platelet endothelial cell adhesion molecule-1 and function in neutrophil transmigration. Mutations in this gene are associated with myeloproliferative diseases. Over-expression of this gene has been found in patients with polycythemia rubra vera. Autoantibodies against the protein may result in pulmonary transfusion reactions, and it may be involved in Wegener's granulomatosis. A related pseudogene, which is adjacent to this gene on chromosome 19, has been identified. [provided by RefSeq, Apr 2014],




Western Blot analysis of K562 cells using CD177 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100



ELK Biotechnology



Western blot analysis of lysate from K562 cells, using CD177 Antibody.