

GDF-5 rabbit pAb

Cat No.: ES4032

For research use only

Overview

Product Name GDF-5 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300.

ELISA: 1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from the Internal region of human

GDF5. AA range:361-410

Specificity GDF-5 Polyclonal Antibody detects endogenous

levels of GDF-5 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Growth/differentiation factor 5

Gene Name GDF5

Cellular localization Secreted . Cell membrane .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 55kD
Human Gene ID 8200
Human Swiss-Prot Number P43026

Alternative Names GDF5; CDMP1; Growth/differentiation factor 5;

GDF-5; Cartilage-derived morphogenetic protein 1;

CDMP-1; Radotermin

Background This gene encodes a secreted ligand of the TGF-beta


(transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that

+86-2


+86-27-59760950 ELKbio@ELKbiotech.com wv



regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. This protein regulates the development of numerous tissue and cell types, including cartilage, joints, brown fat, teeth, and the growth of neuronal axons and dendrites. Mutations in this gene are associated with acromesomelic dysplasia, brachydactyly, chondrodysplasia, multiple synostoses syndrome, proximal symphalangism, and susceptibility to osteoarthritis. [provided by RefSeq, Aug 2016],




Western Blot analysis of 293 cells using GDF-5 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100







Immunohistochemical analysis of paraffin-embedded rat-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-brain, antibody was diluted at 1:100

