

CD294 rabbit pAb

Cat No.: ES4135

For research use only

Overview

Product Name CD294 rabbit pAb

Host species Rabbit

ApplicationsWB;IHC;IF;ELISASpecies Cross-ReactivityHuman;Mouse

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300.

ELISA: 1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from the Internal region of human

PTGDR2. AA range:161-210

Specificity CD294 Polyclonal Antibody detects endogenous

levels of CD294 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Prostaglandin D2 receptor 2

Gene Name PTGDR2

Cell ular localization Cell membrane; Multi-pass membrane protein.

Internalized receptors colocalized with RAB11A. .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 43kD
Human Gene ID 11251
Human Swiss-Prot Number Q9Y5Y4

Alternative Names PTGDR2; CRTH2; DL1R; GPR44; Prostaglandin D2

receptor 2; Chemoattractant receptor-homologous molecule expressed on Th2 cells; G-protein coupled

receptor 44; CD294

Background This gene encodes a G-protein-coupled receptor


that is preferentially expressed in CD4+ effector T helper 2 (Th2) cells. This protein is a prostaglandin




+86-27-59760950 ELKbio@ELKbiotech.com




D2 receptor that mediates the pro-inflammatory chemotaxis of eosinophils, basophils, and Th2 lymphocytes generated during allergic inflammation. Single nucleotide polymorphisms in the 3' UTR of this gene have been associated with asthma susceptibility.[provided by RefSeq, Mar 2011],



Western Blot analysis of 293 cells using CD294 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





Western blot analysis of lysate from 293 cells, using PTGDR2 Antibody.

