

11β-HSD1 rabbit pAb

Cat No.: ES4290

For research use only

Overview

Product Name 11β-HSD1 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300.

ELISA: 1/10000. Not yet tested in other applications.

 $\label{eq:synthesized} \mbox{Immunogen} \qquad \qquad \mbox{Synthesized peptide derived from 11β-HSD1 . at AA}$

range: internal

Specificity 11β-HSD1 Polyclonal Antibody detects endogenous

levels of 11β-HSD1 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Corticosteroid 11-beta-dehydrogenase isozyme 1

Gene Name HSD11B1

Cellular localization Endoplasmic reticulum membrane; Single-pass type

II membrane protein.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 35kD
Human Gene ID 3290
Human Swiss-Prot Number P28845

Alternative Names HSD11B1; HSD11L; Corticosteroid

11-beta-dehydrogenase isozyme 1;

11-beta-hydroxysteroid dehydrogenase 1; 11-DH;

11-beta-HSD1

Background hydroxysteroid 11-beta dehydrogenase 1(HSD11B1)

Homo sapiens The protein encoded by this gene


is a microsomal enzyme that catalyzes the

conversion of the stress hormone cortisol to the






inactive metabolite cortisone. In addition, the encoded protein can catalyze the reverse reaction, the conversion of cortisone to cortisol. Too much cortisol can lead to central obesity, and a particular variation in this gene has been associated with obesity and insulin resistance in children. Mutations in this gene and H6PD (hexose-6-phosphate dehydrogenase (glucose 1-dehydrogenase)) are the cause of cortisone reductase deficiency. Alternate splicing results in multiple transcript variants encoding the same protein.[provided by RefSeq, May 2011],




Western Blot analysis of K562 cells using 11 β -HSD1 Polyclonal Antibody. Antibody was diluted at 1:500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100







Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100

