



IL-12R β 1 rabbit pAb

Cat No.:ES4344

For research use only


Overview

Product Name	IL-12R β 1 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human IL12RB1. AA range:211-260
Specificity	IL-12R β 1 Polyclonal Antibody detects endogenous levels of IL-12R β 1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Interleukin-12 receptor subunit beta-1
Gene Name	IL12RB1
Cellular localization	Membrane; Single-pass type I membrane protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	73kD
Human Gene ID	3594
Human Swiss-Prot Number	P42701
Alternative Names	IL12RB1; IL12R; IL12RB; Interleukin-12 receptor subunit beta-1; IL-12 receptor subunit beta-1; IL-12R subunit beta-1; IL-12R-beta-1; IL-12RB1; IL-12 receptor beta component; CD212
Background	interleukin 12 receptor subunit beta 1(IL12RB1) Homo sapiens The protein encoded by this gene is a type I transmembrane protein that belongs to the hemopoietin receptor superfamily. This protein





binds to interleukine 12 (IL12) with a low affinity, and is thought to be a part of IL12 receptor complex. This protein forms a disulfide-linked oligomer, which is required for its IL12 binding activity. The coexpression of this and IL12RB2 proteins was shown to lead to the formation of high-affinity IL12 binding sites and reconstitution of IL12 dependent signaling. Mutations in this gene impair the development of interleukin-17-producing T lymphocytes and result in increased susceptibility to mycobacterial and Salmonella infections. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014],



Western Blot analysis of HeLa cells using IL-12Rβ1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

