

IL-10Rα rabbit pAb

Cat No.: ES4360

For research use only

Overview

Product Name IL-10Rα rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from the Internal region of human

IL10RA. AA range:151-200

Specificity IL-10Rα Polyclonal Antibody detects endogenous

levels of IL-10Rα protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Interleukin-10 receptor subunit alpha

Gene Name IL10RA

Cell ular localization Cell membrane ; Single-pass type I membrane

protein. Cytoplasm.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 63kD
Human Gene ID 3587
Human Swiss-Prot Number Q13651

Alternative Names IL10RA; IL10R; Interleukin-10 receptor subunit

alpha; IL-10 receptor subunit alpha; IL-10R subunit alpha; IL-10RA; CDw210a; Interleukin-10 receptor

subunit 1; IL-10R subunit 1; IL-10R1; CD210

Background The protein encoded by this gene is a receptor for


interleukin 10. This protein is structurally related to interferon receptors. It has been shown to mediate



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





the immunosuppressive signal of interleukin 10, and thus inhibits the synthesis of proinflammatory cytokines. This receptor is reported to promote survival of progenitor myeloid cells through the insulin receptor substrate-2/PI 3-kinase/AKT pathway. Activation of this receptor leads to tyrosine phosphorylation of JAK1 and TYK2 kinases. Two transcript variants, one protein-coding and the other not protein-coding, have been found for this gene. [provided by RefSeq, Jan 2009],

Western Blot analysis of PC12 cells using IL-10R α Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



+86-27-59760950

ELKbio@ELKbiotech.com