

Olfactory receptor 2F2 rabbit pAb

Cat No.: ES4785

For research use only

Overview

Product Name Olfactory receptor 2F2 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR2F2. AA

range:93-142

Specificity Olfactory receptor 2F2 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 2F2

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 2F2

Gene Name OR2F2

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 32kD
Human Gene ID 135948
Human Swiss-Prot Number O95006

Alternative Names OR2F2; Olfactory receptor 2F2; Olfactory receptor

7-1; OR7-1; Olfactory receptor OR7-6

Background olfactory receptor family 2 subfamily F member


2(OR2F2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor



+86-27-59760950 ELKbio@ELKbiotech.com



proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



+86-27-59760950

Western blot analysis of lysates from NIH/3T3 cells, using OR2F2 Antibody. The lane on the right is blocked with the synthesized peptide.

