

## TFIIA-α rabbit pAb

**Cat No.: ES5685** 

For research use only

## Overview

**Product Name** TFIIA-α rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human TF2A1. AA

range:281-330

**Specificity** TFIIA-α Polyclonal Antibody detects endogenous

levels of TFIIA- $\alpha$  protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Store at -20°C.** Avoid repeated freeze-thaw cycles.

**Protein Name** Transcription initiation factor IIA subunit 1

**Gene Name** GTF2A1 **Cellular localization** Nucleus.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band42kDHuman Gene ID2957Human Swiss-Prot NumberP52655

Alternative Names GTF2A1; TF2A1; Transcription initiation factor IIA

subunit 1; General transcription factor IIA subunit 1; TFIIAL; Transcription initiation factor TFIIA 42 kDa

subunit; TFIIA-42

**Background** Accurate transcription initiation on TATA-containing


class II genes involves the ordered assembly of RNA polymerase II (POLR2A; MIM 180660) and several




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.cor



general initiation factors (summarized by DeJong and Roeder, 1993 [PubMed 8224848]). One of these factors is TFIIA, which when purified from HeLa extracts consists of 35-, 19-, and 12-kD subunits.[supplied by OMIM, Jul 2010],



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by i



Western blot analysis of lysates from Jurkat and HUVEC cells, using TF2A1 Antibody. The lane on the right is blocked with the synthesized peptide.

