



# Hexb rabbit pAb

Cat No.:ES5730

For research use only

## Overview

<b>Product Name</b>	Hexb rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Species Cross-Reactivity</b>	Human;Rat;Mouse;
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human HEXB. AA range:481-530
<b>Specificity</b>	Hexb Polyclonal Antibody detects endogenous levels of Hexb protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	Beta-hexosaminidase subunit beta
<b>Gene Name</b>	HEXB
<b>Cellular localization</b>	Lysosome . Cytoplasmic vesicle, secretory vesicle, Cortical granule .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	63kD
<b>Human Gene ID</b>	3074
<b>Human Swiss-Prot Number</b>	P07686
<b>Alternative Names</b>	HEXB; HCC7; Beta-hexosaminidase subunit beta; Beta-N-acetylhexosaminidase subunit beta; Hexosaminidase subunit B; Cervical cancer proto-oncogene 7 protein; HCC-7; N-acetyl-beta-glucosaminidase subunit beta







## Background

Hexosaminidase B is the beta subunit of the lysosomal enzyme beta-hexosaminidase that, together with the cofactor GM2 activator protein, catalyzes the degradation of the ganglioside GM2, and other molecules containing terminal N-acetyl hexosamines. Beta-hexosaminidase is composed of two subunits, alpha and beta, which are encoded by separate genes. Both beta-hexosaminidase alpha and beta subunits are members of family 20 of glycosyl hydrolases. Mutations in the alpha or beta subunit genes lead to an accumulation of GM2 ganglioside in neurons and neurodegenerative disorders termed the GM2 gangliosidoses. Beta subunit gene mutations lead to Sandhoff disease (GM2-gangliosidosis type II). Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2014],

Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using HEXB Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat cells, using HEXB Antibody. The lane on the right is blocked with the synthesized peptide.

