

IFI-56K rabbit pAb

Cat No.: ES5832

For research use only

Overview

Product Name IFI-56K rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. The antiserum was produced against synthesized

Immunogen The antiserum was produced against synthesized

peptide derived from human IFIT1. AA range:41-90

Specificity IFI-56K Polyclonal Antibody detects endogenous

levels of IFI-56K protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Interferon-induced protein with tetratricopeptide

repeats 1

Gene Name IFIT1

Cellular localization Cytoplasm .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 55kD
Human Gene ID 3434
Human Swiss-Prot Number P09914

Alternative Names IFIT1; G10P1; IFI56; IFNAI1; ISG56;

Interferon-induced protein with tetratricopeptide repeats 1; IFIT-1; Interferon-induced 56 kDa protein;

IFI-56K; P56

Background This gene encodes a protein containing


tetratricopeptide repeats that was originally identified as induced upon treatment with




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.cor



interferon. The encoded protein may inhibit viral replication and translational initiation. This gene is located in a cluster on chromosome 10 with five other closely related genes. There is a pseudogene for this gene on chromosome 13. Alternatively spliced transcript variants encoding multiple isoforms have been observed. [provided by RefSeq, Aug 2012],



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by i



Western blot analysis of lysates from LOVO cells, using IFIT1 Antibody. The lane on the right is blocked with the synthesized peptide.

