

## KV2.1 rabbit pAb

Cat No.:ES5994

For research use only

## Overview

Product Name KV2.1 rabbit pAb

Host species Rabbit

Applications WB;ELISA;IHC Species Cross-Reactivity Human;Mouse;Rat

**Recommended dilutions** WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 **Immunogen** The antiserum was produced against synthesized

peptide derived from human Kv2.1. AA

range:771-820

**Specificity** KV2.1 Polyclonal Antibody detects endogenous

levels of KV2.1 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Potassium voltage-gated channel subfamily B

member 1

Gene Name KCNB1

**Cellular localization** Cell membrane . Perikaryon . Cell projection, axon .

Cell projection, dendrite. Membrane; Multi-pass

membrane protein. Cell junction, synapse,

postsynaptic cell membrane . Cell junction, synapse . Cell junction, synapse, synaptosome . Lateral cell

membra

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 90kD
Human Gene ID 3745
Human Swiss-Prot Number Q14721

Alternative Names KCNB1; Potassium voltage-gated channel subfamily

B member 1; Delayed rectifier potassium channel 1; DRK1; h-DRK1; Voltage-gated potassium channel






**Background** 



Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. Four sequence-related potassium channel genes - shaker, shaw, shab, and shal - have been identified in Drosophila, and each has been shown to have human homolog(s). This gene encodes a member of the potassium channel, voltage-gated, shab-related subfamily. This member is a delayed rectifier potassium channel and its activity is modulated by some other family members. [provided by RefSeq, Jul 2008],



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

