

MYL6 rabbit pAb

Cat No.:ES6313

For research use only

Overview

Product Name	MYL6 rabbit pAb
Host species	Rabbit
Applications	WB;IHC
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	WB 1:500-2000;IHC-p 1:50-300
Immunogen	Synthesized peptide derived from the Internal
	region of human MYL6.
Specificity	MYL6 Polyclonal Antibody detects endogenous
	levels of MYL6 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Myosin light polypeptide 6
Gene Name	MYL6
Cellular localization	cytosol,brush border,membrane,myosin
	complex, unconventional myosin
	complex, extracellular matrix, vesicle, extracellular
	exosome,
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	17kD
Human Gene ID	4637
Human Swiss-Prot Number	P60660
Alternative Names	MYL6; Myosin light polypeptide 6; 17 kDa myosin
	light chain; LC17; Myosin light chain 3; MLC-3;
	Myosin light chain alkali 3; Myosin light chain A3;
	Smooth muscle and nonmuscle myosin light chain
	alkali 6
Background	Myosin is a hexameric ATPase cellular motor
	protein. It is composed of two heavy chains, two



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



nonphosphorylatable alkali light chains, and two phosphorylatable regulatory light chains. This gene encodes a myosin alkali light chain that is expressed in smooth muscle and non-muscle tissues. Genomic sequences representing several pseudogenes have been described and two transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C