

## PDE4B/C/D (phospho Ser133/119/190) rabbit pAb

Cat No.:ES6524

For research use only

## Overview

Product Name PDE4B/C/D (phospho Ser133/119/190) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. The antiserum was produced against synthesize

Immunogen The antiserum was produced against synthesized

peptide derived from human PDE4D around the

phosphorylation site of Ser190/53. AA

range:156-205

Specificity Phospho-PDE4B/C/D (S133/119/190) Polyclonal

Antibody detects endogenous levels of PDE4B/C/D protein only when phosphorylated at \$133/119/190.

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name cAMP-specific 3',5'-cyclic phosphodiesterase 4D

Gene Name PDE4B/PDE4C/PDE4D

Cellular localization[Isoform PDE4B5]: Cytoplasm . Cell membrane .PurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 76kD
Human Gene ID 5144

Human Swiss-Prot Number Q07343/Q08493/Q08499

Alternative Names PDE4B; DPDE4; cAMP-specific 3'; 5'-cyclic

phosphodiesterase 4B; DPDE4; PDE32; PDE4C; DPDE1; cAMP-specific 3',5'-cyclic phosphodiesterase

4C; DPDE1; PDE21; PDE4D; DPDE3; cAMP-specific




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




**Background** 

3',5'-cyclic phosphodiesterase 4D; DPDE3; PDE43
This gene is a member of the type IV, cyclic AMP
(cAMP)-specific, cyclic nucleotide
phosphodiesterase (PDE) family. The encoded
protein regulates the cellular concentrations of
cyclic nucleotides and thereby play a role in signal
transduction. Altered activity of this protein has
been associated with schizophrenia and bipolar
affective disorder. Alternative splicing and the use of
alternative promoters results in multiple transcript
variants encoding different isoforms. [provided by
RefSeq, Jul 2014],




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PDE4D (Phospho-Ser190/53) Antibody



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using PDE4D (Phospho-Ser190/53) Antibody. The picture on the right is blocked with the phospho peptide.







Western blot analysis of lysates from K562 cells treated with H2O2 100uM 30', using PDE4D (Phospho-Ser190/53) Antibody. The lane on the right is blocked with the phospho peptide.



+86-27-59760950