

Pin1 (phospho Ser16) rabbit pAb

Cat No.:ES6594

For research use only

Overview

Product Name	Pin1 (phospho Ser16) rabbit pAb	
Host species	Rabbit	
Applications	WB;IHC;IF;ELISA	
Species Cross-Reactivity	Human;Mouse;Rat;Monkey	
Recommended dilutions	Western Blot: 1/500 - 1/2000.	
	Immunohistochemistry: 1/100 - 1/300. ELISA:	
	1/20000. Not yet tested in other applications.	
Immunogen	The antiserum was produced against synthesized	
_	peptide derived from human Pin1 around the	
	phosphorylation site of Ser16. AA range:1-50	
Specificity	Phospho-Pin1 (S16) Polyclonal Antibody detects	
. ,	endogenous levels of Pin1 protein only when	
	phosphorylated at \$16.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20° C Avoid repeated freeze-thaw cycles	
Protein Name	Pentidyl-prolyl cis-trans isomerase NIMA-interacting	
	1	
Gene Name	PIN1	
Cellular localization	Nucleus Nucleus speckle Cytoplasm Colocalizes	
	with NEK6 in the nucleus (PubMod:16476580)	
	Mainly localized in the nucleus (rubined.10470380).	
	at Cor 71 by DADK1 results in inhibition of its pusher	
	at Set-71 by DAPK1 results in inhibition of its nuclear	
Devification	The entitle during official sources of the source of the s	
Purification	The antibody was aninity-purnied from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	
Observed band	18kD	البر
Human Gene ID	5300	
Human Swiss-Prot Number	Q13526	
Alternative Names	PIN1; Peptidyl-prolyl cis-trans isomerase	



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Background

NIMA-interacting 1; Peptidyl-prolyl cis-trans isomerase Pin1; PPlase Pin1; Rotamase Pin1 Peptidyl-prolyl cis/trans isomerases (PPlases) catalyze the cis/trans isomerization of peptidyl-prolyl peptide bonds. This gene encodes one of the PPIases, which specifically binds to phosphorylated ser/thr-pro motifs to catalytically regulate the post-phosphorylation conformation of its substrates. The conformational regulation catalyzed by this PPIase has a profound impact on key proteins involved in the regulation of cell growth, genotoxic and other stress responses, the immune response, induction and maintenance of pluripotency, germ cell development, neuronal differentiation, and survival. This enzyme also plays a key role in the pathogenesis of Alzheimer's disease and many cancers. Multiple alternatively spliced transcript variants have been found for this gene.[provided by RefSeq, Jun 2011],



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Pin1 (Phospho-Ser16) Antibody

Immunohistochemistry analysis of paraffin-embedded

human heart, using Pin1 (Phospho-Ser16) Antibody. The picture on the right is blocked with the phospho peptide.





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Western blot analysis of lysates from COS7 cells treated with insulin 0.01U/ml 15', using Pin1 (Phospho-Ser16) Antibody. The lane on the right is blocked with the phospho peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C