

SLC6A15 rabbit pAb

Cat No.: ES6709

For research use only

Overview

Product Name SLC6A15 rabbit pAb

Host species Rabbit WB;IHC **Applications**

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions WB 1:500-2000;IHC-p 1:50-300

Immunogen The antiserum was produced against synthesized

peptide derived from human SLC6A15. AA

range:32-81

Specificity SLC6A15 Polyclonal Antibody detects endogenous

levels of SLC6A15 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage Protein Name** Sodium-dependent neutral amino acid transporter

B(0)AT2

SLC6A15 **Gene Name**

Cellular localization Membrane; Multi-pass membrane protein. Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 85kD **Human Gene ID** 55117 **Human Swiss-Prot Number** Q9H2J7

Alternative Names SLC6A15; BOAT2; NTT73; SBAT1; Sodium-dependent

neutral amino acid transporter B(0)AT2; Sodium-

and chloride-dependent neurotransmitter

transporter NTT73; Sodium-coupled branched-chain amino-acid transporter 1; Solute carrier family 6

member

This gene encodes a member of the solute carrier **Background**

family 6 protein family which transports neutral




+86-27-59760950 ELKbio@ELKbiotech.com



JK -- 117 -- 85 -- 48 -- 34 -- 26 -- 19

amino acids. The encoded protein is thought to play a role in neuronal amino acid transport (PMID: 16185194) and may be associated with major depression (PMID: 21521612). Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Feb 2012],

Western blot analysis of lysates from Jurkat cells, using SLC6A15 Antibody. The lane on the right is blocked with the synthesized peptide.



(kD)

Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

