

PKD1 (phospho Ser910) rabbit pAb

Cat No.: ES6785

For research use only

Overview

PKD1 (phospho Ser910) rabbit pAb **Product Name**

Host species Rabbit

WB;IHC;IF;ELISA **Applications Species Cross-Reactivity** Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

The antiserum was produced against synthesized **Immunogen**

peptide derived from human PKD1/PKC mu around

the phosphorylation site of Ser910. AA

range:863-912

Specificity Phospho-PKD1 (S910) Polyclonal Antibody detects

endogenous levels of PKD1 protein only when

phosphorylated at S910.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage**

Serine/threonine-protein kinase D1 **Protein Name**

Gene Name PRKD1

Cellular localization Cytoplasm . Cell membrane . Golgi apparatus,

> trans-Golgi network . Translocation to the cell membrane is required for kinase activation. The antibody was affinity-purified from rabbit

Purification

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 117kD **Human Gene ID** 5587 **Human Swiss-Prot Number** Q15139

Alternative Names PRKD1; PKD; PKD1; PRKCM;

> Serine/threonine-protein kinase D1; Protein kinase C mu type; Protein kinase D; nPKC-D1; nPKC-mu PRKD1 is a serine/threonine kinase that regulates a




Background


+86-27-59760950 ELKbio@ELKbiotech.com




variety of cellular functions, including membrane receptor signaling, transport at the Golgi, protection from oxidative stress at the mitochondria, gene transcription, and regulation of cell shape, motility, and adhesion (summary by Eiseler et al., 2009 [PubMed 19329994]).[supplied by OMIM, Nov 2010],



Western Blot analysis of A431 cells using Phospho-PKD1 (S910) Polyclonal Antibody



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PKD1/PKC mu (Phospho-Ser910) Antibody



Western blot analysis of lysates from A431 cells, using PKD1/PKC mu (Phospho-Ser910) Antibody. The lane on the right is blocked with the phospho peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com





Immunohistochemical analysis of paraffin-embedded human small intestinal carcinoma tissue. 1,primary Antibody was diluted at 1:200(4° overnight). 2, Sodium citrate pH 6.0 was used for antigen retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200



+86-27-59760950