

Rab 2A rabbit pAb

Cat No.:ES6963

For research use only

Overview

Product Name	Rab 2A rabbit pAb	
Host species	Rabbit	
Applications	WB;IHC;IF;ELISA	
Species Cross-Reactivity	Human; Mouse; Rat	
Recommended dilutions	Western Blot: 1/500 - 1/2000.	
	Immunohistochemistry: 1/100 - 1/300. ELISA:	
	1/20000. Not yet tested in other applications.	
Immunogen	The antiserum was produced against synthesized	
	peptide derived from human RAB2A. AA	
	range:84-133	
Specificity	Rab 2A Polyclonal Antibody detects endogenous	
	levels of Rab 2A protein.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.	
Protein Name	Ras-related protein Rab-2A	
Gene Name	RAB2A	
Cellular localization	Endoplasmic reticulum-Golgi intermediate	
	compartment membrane ; Lipid-anchor .	
	Melanosome . Endoplasmic reticulum membrane ;	
	Lipid-anchor . Golgi apparatus membrane ;	
	Lipid-anchor . Cytoplasmic vesicle, secretory vesicle,	
	acrosome . Localized in the Golgi	
Purification	The antibody was affinity-purified from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	
Observed band	25kD	
Human Gene ID	5862	JIIII
Human Swiss-Prot Number	P61019	
Alternative Names	RAB2A; RAB2; Ras-related protein Rab-2A	
Background	The protein encoded by this gene belongs to the	



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Rab family, members of which are small molecular weight guanosine triphosphatases (GTPases) that contain highly conserved domains involved in GTP binding and hydrolysis. The Rabs are membrane-bound proteins, involved in vesicular fusion and trafficking. This protein is a resident of pre-Golgi intermediates, and is required for protein transport from the endoplasmic reticulum (ER) to the Golgi complex. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2011],

Western Blot analysis of A549 cells using Rab 2A Polyclonal Antibody





Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C