

Rab 5C rabbit pAb

Cat No.:ES6965

For research use only

Overview

Purification

Product Name Rab 5C rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human RAB5C. AA

range:122-171

Specificity Rab 5C Polyclonal Antibody detects endogenous

levels of Rab 5C protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Ras-related protein Rab-5C

Gene Name RAB5C

Cellular localization Cell membrane; Lipid-anchor; Cytoplasmic side.

Early endosome membrane; Lipid-anchor.

Melanosome . Identified by mass spectrometry in melanosome fractions from stage I to stage IV. . The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 30kD
Human Gene ID 5878
Human Swiss-Prot Number P51148

Alternative Names RAB5C; RABL; Ras-related protein Rab-5C; L1880;

RAB5L

Background Members of the Rab protein family are small


GTPases of the Ras superfamily that are thought to




+86-27-59760950 ELKbio@ELKbiotech.com www.elkb



ensure fidelity in the process of docking and/or fusion of vesicles with their correct acceptor compartment (Han et al., 1996 [PubMed 8646882]).[supplied by OMIM, Nov 2010],



Western blot analysis of lysates from LOVO, HeLa, and Jurkat cells, using RAB5C Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from 293 cells using RAB5C antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

