

## E2A (phospho Thr355) rabbit pAb

Cat No.: ES7349

For research use only

## Overview

Product Name E2A (phospho Thr355) rabbit pAb

Host species Rabbit
Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

**Recommended dilutions** Immunohistochemistry: 1/100 - 1/300. ELISA:

1/5000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human E2A around the phosphorylation site of Thr355. AA range:321-370

**Specificity** Phospho-E2A (T355) Polyclonal Antibody detects

endogenous levels of E2A protein only when

phosphorylated at T355.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Store at -20°C.** Avoid repeated freeze-thaw cycles.

Protein Name Transcription factor E2-alpha

**Gene Name** TCF3 **Cellular localization** Nucleus .

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

**Observed band** 

Human Gene ID 6929 Human Swiss-Prot Number P15923

Alternative Names TCF3; BHLHB21; E2A; ITF1; Transcription factor

E2-alpha; Class B basic helix-loop-helix protein 21; bHLHb21; Immunoglobulin enhancer-binding factor E12/E47; Immunoglobulin transcription factor 1; Kappa-E2-binding factor; Transcription facto

**Background** This gene encodes a member of the E protein (class


I) family of helix-loop-helix transcription factors. E




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.co



proteins activate transcription by binding to regulatory E-box sequences on target genes as heterodimers or homodimers, and are inhibited by heterodimerization with inhibitor of DNA-binding (class IV) helix-loop-helix proteins. E proteins play a critical role in lymphopoiesis, and the encoded protein is required for B and T lymphocyte development. Deletion of this gene or diminished activity of the encoded protein may play a role in lymphoid malignancies. This gene is also involved in several chromosomal translocations that are associated with lymphoid malignancies including pre-B-cell acute lymphoblastic leukemia (t(1;19), with PBX1), childhood leukemia (t(19;19), with TFPT) and acute leukemia (t(12;19), with ZNF384). Alternatively spliced transcript variants encoding multiple isoforms have bee



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using E2A (Phospho-Thr355) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using E2A (Phospho-Thr355) Antibody. The picture on the right is blocked with the phospho peptide.

