



TSG-6 rabbit pAb

Cat No.:ES7403

For research use only


Overview

Product Name	TSG-6 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human TSG6. AA range:21-70
Specificity	TSG-6 Polyclonal Antibody detects endogenous levels of TSG-6 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Tumor necrosis factor-inducible gene 6 protein
Gene Name	TNFAIP6
Cellular localization	Secreted .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	31kD
Human Gene ID	7130
Human Swiss-Prot Number	P98066
Alternative Names	TNFAIP6; TSG6; Tumor necrosis factor-inducible gene 6 protein; Hyaluronate-binding protein; TNF-stimulated gene 6 protein; TSG-6; Tumor necrosis factor alpha-induced protein 6; TNF alpha-induced protein 6
Background	The protein encoded by this gene is a secretory protein that contains a hyaluronan-binding domain,






and thus is a member of the hyaluronan-binding protein family. The hyaluronan-binding domain is known to be involved in extracellular matrix stability and cell migration. This protein has been shown to form a stable complex with inter-alpha-inhibitor (I alpha I), and thus enhance the serine protease inhibitory activity of I alpha I, which is important in the protease network associated with inflammation. This gene can be induced by proinflammatory cytokines such as tumor necrosis factor alpha and interleukin-1. Enhanced levels of this protein are found in the synovial fluid of patients with osteoarthritis and rheumatoid arthritis.[provided by RefSeq, Dec 2010],



Western blot analysis of lysates from Jurkat, HeLa, and HepG2 cells, using TSG6 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

